



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Solution by J. W. NICHOLSON, A. M., LL. D., President and Professor of Mathematics in the Louisiana State University, and Agricultural and Mechanical College, Baton Rouge, Louisiana.

Let x^2 , $2(n^2-1)x^2$, $2(n^2+1)x^2$, be the three integers. The first condition gives $4n^2x^4 = \square$; hence we have only to solve

$$\left[x^2 \right]^3 + \left[2(n^2-1)x^2 \right]^3 + \left[2(n^2+1)x^2 \right]^3 = \square,$$

$$\text{or } x^6 \left[16n^6 + 48n^2 + 1 \right] = \square,$$

$$\therefore 16n^6 = \left(\frac{48}{2} n^2 \right)^2; \text{ whence } n^2 = 36; \text{ and the three integers are } x^2,$$

$70x^2$ and $74x^2$, where x is any integer.

Also solved by H. W. DRAUGHON, and G. B. M. ZERR.

24. Proposed by F. P. MATZ, M. Sc., Ph. D., Professor of Mathematics and Astronomy in New Windsor College, New Windsor, Maryland.

Solve generally: The sum of the cubes of n consecutive numbers is a square. Determine the numbers, when $n=2$, $n=3$, $n=4$, and $n=5$.

I. Solution by M. A. GRUBER, A. M., War Department, Washington, D. C.

Let m , $m+1$, $m+2$, $m+3$, etc., represent any consecutive numbers the sum of whose cubes is to be taken.

Solving by the differential method, we obtain

$$S = \frac{n}{4} \left\{ n^3 + (4m-2)n^2 + (6m^2-6m+1)n + 4m^3-6m^2+2m \right\}.$$

This reduces to $S = \frac{1}{4} \{ [n(n+2m-1)]^2 + 2m(m-1)[n(n+2m-1)] \}$, (A).

If the sum of the consecutive cubes is to be a square, then

$$[n(n+2m-1)]^2 + 2m(m-1)[n(n+2m-1)] = \square = a^2.$$

Adding $[m(m-1)]^2$ to both members, we have

$$[n(n+2m-1) + m(m-1)]^2 = a^2 + [m(m-1)]^2.$$

This is of the form $(p^2+q^2)^2 = (2pq)^2 + (p^2-q^2)^2$.

Equating the respective values, and reducing for m and n , we obtain

$2m = 1 + \sqrt{4p^2 - 4q^2 + 1}$, and $2n = \sqrt{4p^2 + 4q^2 + 1} - \sqrt{4p^2 - 4q^2 + 1}$. It will be observed that each of the radical quantities is of the form of an odd square, $4p+1$.

There are two conditions that will render the radicals rational, and, at the same time, have a an integer:—

(1) When $4q^2 = 4p$. Then $m = p = q^2$, and $n = 1$. According to this condition there is but *one cube* that can be taken at one time, and hence there would be no *sum of cubes*. This cube is the cube of a square, and is, therefore, also the square of a cube.

(2) The second condition is when $p^2 = q^2$. Then $m = 1$; and substituting this value in (A), we obtain $S = \left\{ \frac{n(n+1)}{2} \right\}^2$, which is the square of the sum of the series, $1+2+3+\dots+n$. From this, then, we have $1^3+2^3+3^3+\dots+n^3 = (1+2+3+\dots+n)^2$, or *the square of the sum of the first n natural numbers is equal to the sum of their respective cubes*.

Therefore, in order that the sum of the cubes of n consecutive numbers be a square, *the first number must be unity.*

When $n=2$, the numbers are 1 and 2; when $n=3$, the numbers are 1, 2, and 3; &c., &c.

Also solved by *Professor COOPER D. SCHMITT, and the PROPOSER.*

II. Solution by B. F. FINKEL, A. M., Professor of Mathematics in Kidder Institute, Kidder, Missouri.

$$\text{Let } S=1+2+3+\dots+n=(n+1)n/2;$$

$$S_2=1^2+2^2+3^2+\dots+n^2=n(n+1)(n+2)/6; \text{ and}$$

$$S_3=1^3+2^3+3^3+\dots+n^3=?$$

$$\begin{aligned} \text{Now } (n+1)^4-n^4 &= 4n^3+6n^2+4n+1 \\ n^4-(n-1)^4 &= 4n^3-6n^2+4n-1=4(n-1)^3+6(n-1)^2+4(n-1)+1 \\ (n-1)^4-(n-2)^4 &= 4n^3-18n^2+28n-15=4(n-2)^3+6(n-2)^2+4(n-2)+1 \\ (n-2)^4-(n-3)^4 &= 4n^3-30n^2+76n-65=4(n-3)^3+6(n-3)^2+4(n-3)+1 \\ &\dots\dots\dots = \dots\dots\dots = \dots\dots\dots \\ 5^4-4^4 &= 369 = 4 \times 4^3 + 6 \times 4^2 + 4 \times 4 + 1 \\ 4^4-3^4 &= 175 = 4 \times 3^3 + 6 \times 3^2 + 4 \times 3 + 1 \\ 3^4-2^4 &= 65 = 4 \times 2^3 + 6 \times 2^2 + 4 \times 2 + 1 \\ 2^4-1^4 &= 15 = 4 \times 1^3 + 6 \times 1^2 + 4 \times 1 + 1 \\ 1^4-0^4 &= 1 = 4 \times 0^3 + 6 \times 0^2 + 4 \times 0 + 1 \end{aligned}$$

$$\text{Adding, } (n+1)^4=4S_3+6S_2+4S+n+1.$$

$$\begin{aligned} \text{Whence, } S_3 &= [(n+1)^4-n-1-6S_2-4S]/4, \\ &= [(n+1)^4-n-n(n+1)(n+2)-2(n+1)n]/4, \\ &= [n^4+2n^3+n^2]/4 = [\tfrac{1}{2}n(n+1)]^2. \end{aligned}$$

If $n=2$, $S_3=9$; if $n=3$, $S_3=36$; if $n=4$, $S_3=100$; if $n=5$, $S_3=225$.

NOTE.—The above method is useful in summing the series: $1^r+2^r+3^r+4^r+\dots+nr$, where r is any integer.

PROBLEMS.

32. Proposed by A. H. BELL, Hillsboro, Illinois.

Decompose into its prime factors the number 549755813889.

33. Proposed by M. A. GRUBER, A. M., War Department, Washington, D. C.

Find three different *sixth powers* whose sum is a square.

[The solution of this problem, if possible, is an answer to the note under the solution of Prob. 16.]